arXiv:1501.03242v2 [math.AT] 6 Oct 2015

CERTAIN HOMOTOPY PROPERTIES RELATED TO $\mathbf{map}(\Sigma^n\mathbb{C}P^2, S^m)$

Jin-ho Lee

Abstract. We compute cohomotopy groups of suspended complex plane $\pi^{n+m}(\Sigma^n\mathbb{C}P^2)$ for $m = 6, 7, 8$. Using these results, we classify path components of the spaces $map(\Sigma^n\mathbb{C}P^2, S^m)$ up to homotopy equivalent. We determine the generalized Gottlieb groups $G_n(\mathbb{C}P^2, S^m)$. Finally, we compute homotopy groups of mapping spaces $map(\Sigma^n\mathbb{C}P^2, S^m; f)$ for all generators $[f]$ of $[\Sigma^n\mathbb{C}P^2, S^m]$ and Gottlieb groups of mapping components containing constant map $map(\Sigma^n\mathbb{C}P^2, S^m; *)$.

1. Introduction

Let X and Y be based topological spaces. A major object of homotopy theory is to study $[X, Y]$, the set of homotopy equivalence classes of based maps. In general, if Y is a co-H-group, $[X, Y]$ has group structure. Let ΣX be the suspension of X. Since every suspended space ΣX is co-group, $[\Sigma X, Y]$ has group structure. If ΣX is a sphere S^n, $[S^n, Y]$ is the n-th homotopy group of Y. On the other hand, $[X, S^n]$ is called the n-th cohomotopy set of X and denoted by $\pi^n(X)$. Moreover, if X is a co-H-group, the cohomotopy set is a group and called n-th cohomotopy group of X. Homotopy groups and cohomotopy groups have been studied by many authors and are a major object in algebraic topology.

Another major object of homotopy theory is to investigate the set of (unbased) maps $f : X \to Y$. We denote $map(X, Y)$ to be the set of all continuous maps from X to Y equipped with compact-open topology. Then we write $map(X, Y; f)$ for the path-component of $map(X, Y)$ containing a map $f : X \to Y$. Important cases are $map(X, Y; *)$ and $map(X, X; 1)$ where $* : X \to Y$ is the constant map and $1 : X \to X$ is the identity map on X. D. Harris proved that every topological space X is the space of path components of a stratifiable space $S(X)$ [9]. Thus it is important to study path components of mapping spaces to analyze topological spaces.

Lang proved that if X is a suspended space, then all path components of based mapping space $map_*(X, Y)$ have the same homotopy type [15, Theorem 2.1]. Whitehead proved that $map(S^n, S^m; f)$ is homotopy equivalent to

---

2010 *Mathematics Subject Classification.* Primary 54C35,55P15,55Q52,55Q55.

Key words and phrases. Complex plane, Cohomotopy group, Toda Bracket, mapping space, path-component.

$map(S^n, S^m; *)$ if and only if w_f has a section, where $*: S^n \to S^m$ is a constant map [30, Theorem 2.8]. Smith proved a rational homotopy equivalence

$$map(X, X; *) \simeq_{\mathbb{Q}} X \times map_*(X, X : *)$$

for a "nice" space X [26]. Lupton and Smith proved that

$$map(X, Y; f) \simeq map(X, Y; f + d)$$

for a CW co-H-space X and any CW complex Y, where $d : X \to Y$ is a cyclic map [16, Theorem 3.10]. Recently, Gatsinzi [3] proved that the dimension of the rational Gottlieb group of the universal cover $\widetilde{map}(X, S^{2n}; f)$ of the function space $map(X, S^{2n}; f)$ is at least equal to the dimension of $\tilde{H}^*(X; \mathbb{Q})$ under several assumptions. Lupton and Smith [16] showed that

$$G_n(map_*(X, Y; *)) \cong G_n(Y) \oplus G_n(X, Y).$$

Maruyama and Oshima [18] determined homotopy groups of $map_*(SU(3), SU(3); *)$, $map_*(Sp(2), Sp(2); *)$ and $map_*(G_2, G_2; *)$.

In Section 2, we present some basic knowledge of composition methods [28]. We review a mapping cone sequence and a Puppe sequence related to the suspended complex projective plane and discuss the concept of cyclic maps and its properties. Also we recall the Toda bracket and its properties.

In Sections 3, 4 and 5, we compute $\pi^n(\Sigma^{n+k}\mathbb{C}P^2)$ for $k = 6, 7$. As a result, we obtain the following results:

case n	2	3	4	5	6	7
$\pi^n(\Sigma^{n+6}\mathbb{C}P^2)$	$2+15$	$2+3$	$8+2+3^2+5$	$4+9$	4^2+9+3	$4+3$
case n	8	9	10	11	$n \geq 12$	
$\pi^n(\Sigma^{n+6}\mathbb{C}P^2)$	4^2+3^2	$4+3$	$2+3$	$2+3$	2	

case n	2	3	4	5
$\pi^n(\Sigma^{n+7}\mathbb{C}P^2)$	$2+3$	2^2+21	$4+2^3+21$	$4+2^2+63$
case n	6	7	8	9
$\pi^n(\Sigma^{n+7}\mathbb{C}P^2)$	$4+2^2+63$	$8+2^2$	$8+2^2$	$8+2^2$
case n	10	11	12	$n \geq 13$
$\pi^n(\Sigma^{n+7}\mathbb{C}P^2)$	$\infty+8+2+63$	$8+2+63$	$\infty+8+2+63$	$8+2+63$

case n	2	3	4	5	6	7	8
$\pi^n(\Sigma^{n+8}\mathbb{C}P^2)$	2^2+21	$2+3$	$8+2$	$4+9$	$8+4+9+3$	$8+3$	8^2+2^2
case n	9	10	11	12	13	$n \geq 14$	
$\pi^n(\Sigma^{n+8}\mathbb{C}P^2)$	$8+3$	$8+2$	2^2	$2+3$	2+3	3	

where an integer n denotes the cyclic group $\mathbb{Z}_n$, $(s)^k$ denotes the k-times direct sum of $\mathbb{Z}_s$, ∞ denotes the group of integers $\mathbb{Z}$ and "+" denotes the direct sum of abelian groups.

In section 6, using our result in section 3, 4 and 5, we determine nth-homotopy groups of function spaces $map_*(\mathbb{C}P^2, \mathbb{C}P^2; *)$ for $4 \leq n \leq 13$.

In Section 7, we apply our results to the classification of path components of mapping spaces up to homotopy equivalent and evaluation fibrations up to fibre homotopy equivalent. Hansen proved that the evaluation fibration $w_f : map(\Sigma X, \Sigma Y; f) \to \Sigma Y$ has a section if and only if $[f, id_{\Sigma Y}] = 0$, where $[\ ,\ ]$ is the generalized Whitehead product [8]. Lupton and Smith proved that the following statements are all equivalent: (1) a map $f : X \to Y$ is cyclic, (2) w_f has a section, and (3) two fibrations w_f and w_* are fiber homotopy equivalent, where $*$ is a constant map [16]. Moreover, we apply our results to the formulation of generalized Gottlieb groups from suspended complex plane to sphere and Gottlieb groups of path components of constant map. We use the notation of [28, 11] freely.

2. Preliminaries

The complex projective plane $\mathbb{C}P^2$ is defined by the mapping cone $S^2 \cup_{\eta_2} e^4$ where $\eta_2 : S^3 \to S^2$ is the Hopf fibration. We denote $\eta_k = \Sigma^{k-2}\eta_2$ for $k \geq 2$. Consider a Puppe sequence

$$S^3 \xrightarrow{\eta_2} S^2 \xrightarrow{i} \mathbb{C}P^2 \xrightarrow{p} S^4 \xrightarrow{\eta_3} S^3 \xrightarrow{\Sigma i} \cdots ,$$

where $i : S^2 \to \mathbb{C}P^2$ is the inclusion map and $p : \mathbb{C}P^2 \to S^4$ is the collapsing map of S^2 to a point. The Puppe sequence induces a long exact sequence of homotopy sets

$$\cdots \to \pi_{n+3}(S^m) \xrightarrow{\eta_{n+3}^*} \pi_{n+4}(S^m) \xrightarrow{\Sigma^n p^*} [\Sigma^n\mathbb{C}P^2, S^m]$$
$$\xrightarrow{\Sigma^n i^*} \pi_{n+2}(S^m) \xrightarrow{\eta_{n+2}^*} \pi_{n+3}(S^m) \to \cdots .$$

Then we have the short exact sequence

$$0 \to Coker\eta_{n+3}^* \xrightarrow{\Sigma^n p^*} [\Sigma^n\mathbb{C}P^2, S^m] \xrightarrow{\Sigma^n i^*} Ker\eta_{n+2}^* \to 0. \tag{1.1}$$

Let $g_6(\mathbb{C}) : \Sigma^7\mathbb{C}P^2 \to S^6$ be the S^1-transfer map [23]. This map is the adjoint of the composite of inclusions

$$\Sigma\mathbb{C}P^2 \hookrightarrow SU(3) \hookrightarrow SO(6) \hookrightarrow \Omega^6 S^6.$$

We set $g_{n+6}(\mathbb{C}) = \Sigma^n g_6(\mathbb{C})$ for $n \geq 1$. Note that $g_6(\mathbb{C}) = [\iota_6, \iota_6] \circ \Sigma^7 p + \nu_6 \circ \overline{2\iota_9}$ and $2g_n(\mathbb{C}) = \nu_n \circ \overline{2\iota_{n+3}}$ for $n \geq 7$ [11, Proposition 3.3].

Let G be an abelian group. For a prime p, we denote the p-primary parts of G by $G_{(p)}$. For an odd prime p, we have an isomorphism

$$[\Sigma^n \mathbb{C}P^2, S^k]_{(p)} \cong \pi_{n+2}(S^k)_{(p)} \oplus \pi_{n+4}(S^k)_{(p)}, \tag{1.2}$$

since $\pi_{n+1}(S^n)$ is of order 2 for $n \geq 3$ [28, Proposition 5.1].

The space $\Sigma\mathbb{C}P^{n-1}$ has reduced homology

$$\widetilde{H}_*(\Sigma\mathbb{C}P^{n-1}) = \mathbb{Z}_p[x_3, x_5, \cdots, x_{2n-1}].$$

There is a Steenrod operation on $\widetilde{H}_*(\Sigma\mathbb{C}P^{n-1})$ given by

$$\mathcal{P}^j(x_{2r+1}) = r!/(j! \cdot (r-j)!) \cdot x_{2r+jp+1} \quad (2 \leq r \leq n-1).$$

It has been shown that there exists a wedge decomposition [22]

$$\Sigma\mathbb{C}P^{n-1} \simeq \bigvee_{i=1}^{p-1} A_i(n)$$

where $A_i(n)$ has a reduced homology

$$\widetilde{H}_*(A_i(n)) = \mathbb{Z}/p\mathbb{Z}\{x_{2i+1}, x_{2i+q+1}, \cdots, x_{2i+(r_i^n-1)q+1} \mid r_i^n = \lfloor (n-i-1)/(p-1) \rfloor + 1\}$$

that inherits a Steenrod operation from $\widetilde{H}_*(\Sigma\mathbb{C}P^{n-1})$.

Consider elements $\alpha \in [Y, Z]$, $\beta \in [X, Y]$, and $\gamma \in [W, X]$ which satisfy $\alpha \circ \beta = 0$ and $\beta \circ \gamma = 0$. Let C_β be the mapping cone of β, $i : Y \to C_\beta$ and $p : C_\gamma \to \Sigma X$ be the inclusion and the shrinking map, respectively. We denote an *extension* $\overline{\alpha} \in [C_\beta, Z]$ of α satisfying $i^*(\overline{\alpha}) = \alpha$ and a *coextension* $\widetilde{\gamma} \in [\Sigma W, C_\beta]$ of γ satisfying $p_*(\widetilde{\gamma}) = \Sigma\gamma$ [28]. From the definition of extension, an extension $\overline{\alpha}$ exists if and only if $\alpha \circ \beta = 0$. Similarly, a coextension $\widetilde{\gamma}$ exists if and only if $\beta \circ \gamma = 0$.

We recall a useful relation between extensions and Toda brackets [28, Proposition 1.9]:

Proposition 1. *Let $\alpha \in [Y, Z]$, $\beta \in [X, Y]$, and $\gamma \in [W, X]$ be elements which satisfy $\alpha \circ \beta = 0$ and $\beta \circ \gamma = 0$. Let $\{\alpha, \beta, \gamma\}$ be the Toda bracket and $p : C_\gamma \to \Sigma W$ be the shrinking map. Then, we have*

$$\alpha \circ \overline{\beta} \in \{\alpha, \beta, \gamma\} \circ p.$$

Here, we recall the concept of a cyclic map and Gottlieb group $G_n(X)$ of a space X [6, 29].

Definition 1. *A map $f : Y \to X$ is cyclic if there is a map $F : X \times Y \to X$, called an affiliated map of f, such that the diagram homotopy commutative*

$$\begin{tikzcd}
X \times Y \arrow[r, "F"] & X \\
X \vee Y \arrow[u, "i"] \arrow[r, "1\vee f"] & X \vee X. \arrow[u, "\nabla"']
\end{tikzcd}$$

Let $G(Y, X)$ denote the set of all homotopy classes of cyclic maps from Y to X. Varadarajan showed that $G(Y, X)$ has group structure for any co-H-space Y [29]. For an integer $n \geq 1$, the set of homotopy classes of cyclic maps $\Sigma^n X \to Y$ we denote by $G_n(X, Y)$, and call the *n-th generalized Gottlieb group of* (X, Y). When $Y = S^n$, $G(Y, X) = G_n(X)$ is the *n-th Gottlieb group of* X. In [6], Gottlieb introduced and studied the evaluation subgroups

$$G_n(X) = w_*(\pi_n(map(X, X; 1)))$$

where $w : map(X, X; 1) \to X$ is the evaluation map. Note that the $G_n(X)$ can alternatively be described as homotopy classes of maps $f : S^n \to X$ such that $(f|1) : S^n \vee X \to X$ admits an extension $F : S^n \times X \to X$ up to homotopy.

3. Determination of $[\Sigma^{n+6}\mathbb{C}P^2, S^n]$

In this section, we shall determine the generators of the 2-primary components of n-th cohomotopy groups of $(n+6)$-fold suspended complex projective planes. We analyze the following extension

$$0 \to Coker\,\eta^*_{n+9} \xrightarrow{\Sigma^n p^*} [\Sigma^{n+6}\mathbb{C}P^2, S^n] \xrightarrow{\Sigma^n i^*} Ker\,\eta^*_{n+8} \to 0 \tag{2.1}$$

induced by (1.1).

Lemma 1. *(1) For the homomorphisms $\eta^*_{n+9} : \pi^n_{n+9} \to \pi^n_{10}$, we have the following table of the cokernel of η^*_{n+9} :*

$n =$	*2*	*3*	*4*	*5, 6, 7*	*8*	*9*	*10*	*11*	≥ 12
$Coker\,\eta^*_{n+9}$	$\mathbb{Z}_2$	Z_2	$\mathbb{Z}_8 \oplus \mathbb{Z}_2$	$\mathbb{Z}_4$	$\mathbb{Z}_4^2$	Z_4	$\mathbb{Z}_2$	$\mathbb{Z}_2$	*0*
generators	$\eta_2\mu_3$	ε'	$\nu_4\sigma'$ $\Sigma\varepsilon'$	$\nu_n\sigma_{n+3}$	$\sigma_8\nu_{15}$ $\nu_8\sigma_{11}$	$\sigma_9\nu_{16}$	$\sigma_{10}\nu_{17}$	$\sigma_{11}\nu_{18}$	*0* *0*

(2) The homomorphisms $\eta^n_{n+8} : \pi^n_{n+8} \to \pi^n_{n+9}$ are monomorphic except $n = 6$ *where*

$$Ker\,\eta^*_{14} = \mathbb{Z}_4\{2\bar{\nu}_6\}.$$

From (2.1) and Lemma 1, we state our result.

Proposition 2. *(1)* $[\Sigma^8\mathbb{C}P^2, S^2] = \mathbb{Z}_2\{\eta_2 \circ \mu_3 \circ \Sigma^8 p\} \oplus \mathbb{Z}_{15}$.
(2) $[\Sigma^9\mathbb{C}P^2, S^3] = \mathbb{Z}_2\{\varepsilon' \circ \Sigma^9 p\} \oplus \mathbb{Z}_3$.
(3) $[\Sigma^{10}\mathbb{C}P^2, S^4] = \mathbb{Z}_8\{\nu_4 \circ \sigma' \circ \Sigma^{10} p\} \oplus \mathbb{Z}_2\{\Sigma\varepsilon' \circ \Sigma^{10} p\} \oplus \mathbb{Z}_3^2 \oplus \mathbb{Z}_5$.
(4) $[\Sigma^{11}\mathbb{C}P^2, S^5] = \mathbb{Z}_4\{\nu_5 \circ \sigma_8 \circ \Sigma^{11} p\} \oplus \mathbb{Z}_9$.
(5) $[\Sigma^{12}\mathbb{C}P^2, S^6] = \mathbb{Z}_4^2\{\nu_6 \circ \sigma_9 \circ \Sigma^{12} p, \bar{\nu}_6 \circ \overline{2\iota_{14}}\} \oplus \mathbb{Z}_9 \oplus \mathbb{Z}_3$.
(6) $[\Sigma^{13}\mathbb{C}P^2, S^7] = \mathbb{Z}_4\{\nu_7 \circ \sigma_{10} \circ \Sigma^{13} p\} \oplus \mathbb{Z}_3$.
(7) $[\Sigma^{14}\mathbb{C}P^2, S^8] = \mathbb{Z}_4^2\{\sigma_8 \circ \nu_{15} \circ \Sigma^{14} p, \nu_8 \circ \sigma_{11} \circ \Sigma^{14} p\} \oplus \mathbb{Z}_3^2$.
(8) $[\Sigma^{15}\mathbb{C}P^2, S^9] = \mathbb{Z}_4\{\sigma_9 \circ \nu_{16} \circ \Sigma^{15} p\} \oplus \mathbb{Z}_3$.
(9) $[\Sigma^{16}\mathbb{C}P^2, S^{10}] = \mathbb{Z}_2\{\sigma_{10} \circ \nu_{17} \circ \Sigma^{16} p\} \oplus \mathbb{Z}_3$.
(10) $[\Sigma^{17}\mathbb{C}P^2, S^{11}] = \mathbb{Z}_2\{\sigma_{11} \circ \nu_{18} \circ \Sigma^{17} p\} \oplus \mathbb{Z}_3$.
(11) $[\Sigma^{n+6}\mathbb{C}P^2, S^n] = \mathbb{Z}_3$ *for* $n \geq 12$.

Proof. By Lemma 1 (2), we determine all group structures of $[\Sigma^{n+6}\mathbb{C}P^2, S^n]$ except $n = 6$ so we consider $[\Sigma^{12}\mathbb{C}P^2, S^6]$. By (2.1) and Lemma 1, we have the following commutative ladder:

$$\begin{tikzcd}
0 \arrow[r] & \mathbb{Z}_4\{\nu_6 \circ \sigma_9\} \arrow[r, "\Sigma^{12}p^*"] \arrow[d, "\Sigma_1"] & {[\Sigma^{12}\mathbb{C}P^2, S^6]_{(2)}} \arrow[r, "\Sigma^{12}i^*"] \arrow[d, "\Sigma_2"] & \mathbb{Z}_4\{2\overline{\nu}_6\} \arrow[r] \arrow[d, "\Sigma_3"] & 0 \\
0 \arrow[r] & \mathbb{Z}_4\{\nu_7 \circ \sigma_{10}\} \arrow[r, "\Sigma^{13}p^*"] & {[\Sigma^{13}\mathbb{C}P^2, S^7]_{(2)}} \arrow[r, "\Sigma^{12}i^*"] & 0 \arrow[r] & 0
\end{tikzcd}$$

Since Σ_1 and $\Sigma^{13}p^*$ are isomorphisms, $\Sigma^{12}p^*$ has left inverse. This implies that the first row splits. This completes the proof. ☐

For odd prime p, we have an isomorphism

$$[\Sigma^{n+6}\mathbb{C}P^2, S^n]_{(p)} \cong \pi_{n+8}(S^n)_{(p)} \oplus \pi_{n+10}(S^n)_{(p)} \tag{2.2}$$

by (2.2). From [28, Chapter 13], then, we have the following:

Proposition 3. *The odd primary components of* $[\Sigma^{n+6}\mathbb{C}P^2, S^n]$ *are*
(1) $[\Sigma^8\mathbb{C}P^2, S^2] = \mathbb{Z}_3\{\eta_2 \circ \overline{\alpha_2(3)}\} \oplus \mathbb{Z}_5\{\eta_2 \circ \overline{\alpha_{1,5}(3)}\}$.
(2) $[\Sigma^9\mathbb{C}P^2, S^3] = \mathbb{Z}_3\{\alpha_1(3) \circ \alpha_2(6) \circ \Sigma^9 p\}$.
(3) $[\Sigma^{10}\mathbb{C}P^2, S^4] = \mathbb{Z}_3^2\{\alpha_1(4) \circ \alpha_2(7) \circ \Sigma^{10} p, \nu_4 \circ \alpha_2(7) \circ \Sigma^{10} p\}$.
(4) $[\Sigma^{11}\mathbb{C}P^2, S^5] = \mathbb{Z}_9\{\beta_1(5) \circ \Sigma^{11} p\}$.
(5) $[\Sigma^{12}\mathbb{C}P^2, S^6] = \mathbb{Z}_9\{\beta_1(6) \circ \Sigma^{12} p\} \oplus \mathbb{Z}_3\{[\iota_6, \iota_6] \circ \overline{\alpha_1(11)}\}$.
(6) $[\Sigma^{13}\mathbb{C}P^2, S^7] = \mathbb{Z}_3\{\beta_1(7) \circ \Sigma^{13} p\}$.
(7) $[\Sigma^{14}\mathbb{C}P^2, S^8] = \mathbb{Z}_3^2\{\beta_1(8) \circ \Sigma^{14} p, [\iota_8, \iota_8] \circ \alpha_1(15) \circ \Sigma^{14}\}$.
(8) $[\Sigma^{n+6}\mathbb{C}P^2, S^n] = \mathbb{Z}_3\{\beta_1(n) \circ \Sigma^{n+6} p\}$ *for* $n \geq 8$.

4. Determination of $[\Sigma^{n+7}\mathbb{C}P^2, S^n]$

In this section, we shall determine the generators of the 2-primary components of n-th cohomotopy groups of $(n+7)$-fold suspended complex projective planes. We analyze the short exact sequence

$$0 \to Coker\, \eta_{n+10}^* \xrightarrow{\Sigma^n p^*} [\Sigma^{n+7}\mathbb{C}P^2, S^n] \xrightarrow{\Sigma^n i^*} Ker\, \eta_{n+9}^* \to 0 \tag{3.1}$$

induced by (1.1).

Lemma 2. *(1) For the homomorphisms* $\eta_{n+10}^* : \pi_{n+10}^n \to \pi_{11}^n$*, we have the following table of the cokernel of* η_{n+10}^* *:*

$n =$	*2*	*3*	*4*	*5*	*6, 7, 8, 9*	*10, 11*	*12*	≥ 13
Coker η_{n+10}^*	$\mathbb{Z}_2$	Z_2^2	$\mathbb{Z}_2^4$	$\mathbb{Z}_4 \oplus \mathbb{Z}_2$	$\mathbb{Z}_4 \oplus \mathbb{Z}_2$	$\mathbb{Z}_4$	$\mathbb{Z} \oplus \mathbb{Z}_4$	$\mathbb{Z}_4$
generators	$\eta_2\varepsilon'$	μ' $\varepsilon_3\nu_{11}$	$\Sigma\mu', \nu_4\bar{\nu}_7$ $\nu_4\varepsilon_7, \varepsilon_4\nu_{12}$	ζ_5 $\nu_5\bar{\nu}_8$	ζ_n $\bar{\nu}_n\nu_{n+8}$	ζ_n	$P(\iota_{25})$ ζ_{12}	ζ_n

(2) For the homomorphisms $\eta_{n+9}^* : \pi_{n+9}^n \to \pi_{10}^n$*, we have the following table of the kernel of* η_{n+9}^* *:*

$n =$	*2, 3*	*4, 5, 6*	*7*	*8*	*9*	*10*	≥ 11
Ker η_{n+9}^*	*0*	$\mathbb{Z}_2$	$\mathbb{Z}_2^2$	$\mathbb{Z}_2^2$	$\mathbb{Z}_2^2$	$\mathbb{Z} \oplus \mathbb{Z}_2^2$	$\mathbb{Z}_2^2$
generators	*0*	ν_n^3	ν_7^3 $\sigma'\eta_{14}^2 + \eta_7\varepsilon_8$	ν_8^3 $(E\sigma')\eta_{15}^2 + \eta_8\varepsilon_9$	ν_9^3 $\eta_9\varepsilon_{10}$	$2P(\iota_{21})$ $\nu_{11}^3, \eta_{11}\varepsilon_{12}$	ν_n^3 $\eta_n\varepsilon_{n+1}$

By Lemma 2, we obtain

$$\Sigma^n p^* : Coker\, \eta_{n+10}^* \to [\Sigma^{n+7}\mathbb{C}P^2, S^n]$$

are isomorphic for $n = 2, 3$. So, we have the following:

Proposition 4. *(1)* $[\Sigma^9\mathbb{C}P^2, S^2] = \mathbb{Z}_2\{\eta_2 \circ \varepsilon' \circ \Sigma^9 p\} \oplus \mathbb{Z}_3$.
(2) $[\Sigma^{10}\mathbb{C}P^2, S^3] = \mathbb{Z}_2^2\{\mu' \circ \Sigma^{10}p, \varepsilon_3 \circ \nu_{11} \circ \Sigma^{10}p\} \oplus \mathbb{Z}_3 \oplus \mathbb{Z}_7$.

Proposition 5. $[\Sigma^{11}\mathbb{C}P^2, S^4] = \mathbb{Z}_4\{\nu_4^2 \circ g_{10}(\mathbb{C})\} \oplus \mathbb{Z}_2^3\{(\Sigma\mu') \circ \Sigma^{11}p, \varepsilon_4 \circ \nu_{12} \circ \Sigma^{11}p, \nu_4 \circ \overline{\nu}_7 \circ \Sigma^{11}p\} \oplus \mathbb{Z}_{21}$.
Relation : $\nu_4 \circ \varepsilon_7 \circ \Sigma^{11}p = 2\nu_4^2 \circ g_{10}(\mathbb{C})$.

Proof. By (3.1) and Lemma 2, we have

$$0 \to \mathbb{Z}_2^4\{\Sigma\mu', \nu_4\circ\overline{\nu}_7, \nu_4\circ\varepsilon_7, \varepsilon_4\circ\nu_{12}\}\oplus\mathbb{Z}_{15} \xrightarrow{\Sigma^{11}p^*} [\Sigma^{11}\mathbb{C}P^2, S^4] \xrightarrow{\Sigma^{11}i^*} \mathbb{Z}_2\{\nu_4^3\} \to 0.$$

By [11, Proposition 3.6], we have a relation

$$2\nu_4^2 \circ g_{10}(\mathbb{C}) = \nu_4^3 \circ \overline{2\iota_{13}} = \nu_4 \circ \varepsilon_7 \circ \Sigma^{11}p.$$

This completes the proof. □

Proposition 6. $[\Sigma^{12}\mathbb{C}P^2, S^5] = \mathbb{Z}_4\{\zeta_5\circ\Sigma^{12}p\}\oplus\mathbb{Z}_2^2\{\nu_5\circ\overline{\nu}_8\circ\Sigma^{12}p, \nu_5^2\circ g_{11}(\mathbb{C})\}\oplus \mathbb{Z}_7 \oplus \mathbb{Z}_9$.

Proof. By (3.1) and Lemma 2, we have

$$0 \to \mathbb{Z}_4\{\zeta_5\} \oplus \mathbb{Z}_2\{\nu_5 \circ \overline{\nu}_8\} \oplus \mathbb{Z}_7 \oplus \mathbb{Z}_9 \xrightarrow{\Sigma^{12}p^*} [\Sigma^{12}\mathbb{C}P^2, S^5] \xrightarrow{\Sigma^{12}i^*} \mathbb{Z}_2\{\nu_5^3\} \to 0$$

Since ν_5^2 is of order 2, $\nu_5^2 \circ g_{11}(\mathbb{C}) \in [\Sigma^{12}\mathbb{C}P^2, S^5]$ is of order 2. This completes the proof. □

Proposition 7. $[\Sigma^{13}\mathbb{C}P^2, S^6] = \mathbb{Z}_4\{\zeta_6 \circ \Sigma^{13}p\} \oplus \mathbb{Z}_2^2\{\overline{\nu}_6 \circ \nu_{14} \circ \Sigma^{13}p, \nu_6^2 \circ g_{12}(\mathbb{C})\} \oplus \mathbb{Z}_7 \oplus \mathbb{Z}_9$.

Proof. By (3.1) and Lemma 2, we have

$$0 \to \mathbb{Z}_4\{\zeta_6\} \oplus \mathbb{Z}_2\{\overline{\nu}_6 \circ \nu_{14}\} \oplus \mathbb{Z}_7 \oplus \mathbb{Z}_9 \xrightarrow{\Sigma^{13}p^*} [\Sigma^{13}\mathbb{C}P^2, S^6] \xrightarrow{\Sigma^{13}i^*} \mathbb{Z}_2\{\nu_6^3\} \to 0.$$

Then we have a commutative diagram

$$\begin{tikzcd}
0 \arrow[r] & \mathbb{Z}_4\{\zeta_5\} \oplus \mathbb{Z}_2\{\nu_5 \circ \overline{\nu}_8\} \arrow[r, "\Sigma^{12}p^*"] \arrow[d, "\Sigma_1"] & {[\Sigma^{12}\mathbb{C}P^2, S^5]} \arrow[r, "\Sigma^{12}i^*"] \arrow[d, "\Sigma_2"'] & \mathbb{Z}_2\{\nu_5^3\} \arrow[r] \arrow[d, "\Sigma_3"] & 0 \\
0 \arrow[r] & \mathbb{Z}_4\{\zeta_6\} \oplus \mathbb{Z}_2\{\overline{\nu}_6 \circ \nu_{14}\} \arrow[r, "\Sigma^{13}p^*"] & {[\Sigma^{13}\mathbb{C}P^2, S^6]} \arrow[r, "\Sigma^{13}i^*"] & \mathbb{Z}_2\{\nu_6^3\} \arrow[r] & 0
\end{tikzcd}$$

Since the first row is split and Σ_3 is an isomorphism, the second row is also split. This completes the proof. □

Note that the Hopf map $\sigma_8 : S^{15} \to S^8$ induces an isomorphism

$$[X, S^7] \oplus [\Sigma X, S^{15}] \to [\Sigma X, S^8], \quad (\alpha, \beta) \mapsto \Sigma\alpha + \sigma_8 \circ \beta. \tag{1}$$

Lemma 3.

$$[\Sigma^{14}\mathbb{C}P^2, S^7]_{(2)} \cong [\Sigma^{15}\mathbb{C}P^2, S^8]_{(2)} \cong [\Sigma^{16}\mathbb{C}P^2, S^9]_{(2)}.$$

Proof. By (2) and [11, Proposition 3.2], we have an isomorphism

$$\Sigma : [\Sigma^{14}\mathbb{C}P^2, S^7] \to [\Sigma^{15}\mathbb{C}P^2, S^8].$$

We consider a 2-primary EHP sequence

$$[\Sigma^{17}\mathbb{C}P^2, S^{17}]_{(2)} \xrightarrow{\triangle} [\Sigma^{15}\mathbb{C}P^2, S^8]_{(2)} \xrightarrow{\Sigma} [\Sigma^{16}\mathbb{C}P^2, S^9]_{(2)} \xrightarrow{H}$$

$$\xrightarrow{H} [\Sigma^{16}\mathbb{C}P^2, S^{17}]_{(2)} \xrightarrow{\triangle} [\Sigma^{14}\mathbb{C}P^2, S^8]_{(2)}$$

where $[\Sigma^{17}\mathbb{C}P^2, S^{17}] = 0$, $[\Sigma^{16}\mathbb{C}P^2, S^{17}]_{(2)} = \mathbb{Z}_4\{\nu_{17} \circ \Sigma^{16}p\}$ and $[\Sigma^{14}\mathbb{C}P^2, S^8]_{(2)} = \mathbb{Z}_4^2\{\sigma_8 \circ \nu_{15} \circ \Sigma^{14}p, \nu_8 \circ \sigma_{11} \circ \Sigma^{14}p\}$ by Proposition 3 and [11, Proposition 3.1, 3.2]. Then $\Sigma : [\Sigma^{15}\mathbb{C}P^2, S^8]_{(2)} \to [\Sigma^{16}\mathbb{C}P^2, S^9]_{(2)}$ is injective. By [28, (7.19)], we have

$$\Delta(\nu_{17} \circ \Sigma^{16}p) = P(\nu_{17}) \circ \Sigma^{14}p = 2\sigma_8 \circ \nu_{15} \circ \Sigma^{14}p - x\nu_8 \circ \sigma_{11} \circ \Sigma^{14}p$$

for an odd x. Then $\triangle : [\Sigma^{16}\mathbb{C}P^2, S^{17}]_{(2)} \to [\Sigma^{14}\mathbb{C}P^2, S^8]_{(2)}$ is injective, so that $\Sigma : [\Sigma^{15}\mathbb{C}P^2, S^8]_{(2)} \to [\Sigma^{16}\mathbb{C}P^2, S^9]_{(2)}$ is surjective. Therefore Σ is isomorphic. This completes the proof. □

By [28, (5.5), (7.10), (7.19)], we have

$$\sigma' \circ \eta_{14}^3 = \sigma' \circ 4\nu_{14} = 4(\nu_7 \circ \sigma_{10}) = \eta_7^2 \circ \varepsilon_9 = \eta_7 \circ \varepsilon_8 \circ \eta_{16}.$$

This implies $\sigma' \circ \eta_{14}^3 + \eta_7 \circ \varepsilon_8 \circ \eta_{16} = 0$. Then we can take an *extension* $\overline{\sigma' \circ \eta_{14}^2 + \eta_7 \circ \varepsilon_8} \in [\Sigma^{14}\mathbb{C}P^2, S^7]$ of $\sigma' \circ \eta_{14}^2 + \eta_7 \circ \varepsilon_8$. Similarly, we can take an *extension* $\overline{(E\sigma') \circ \eta_{15}^2 + \eta_8 \circ \varepsilon_9} \in [\Sigma^{15}\mathbb{C}P^2, S^8]$ of $(E\sigma') \circ \eta_{15}^2 + \eta_8 \circ \varepsilon_9$. By [28, (7.5), (7.10), (7.19), Theorem 7.2], we have

$$\eta_{n+9}^*(\eta_n \varepsilon_{n+1}) = \eta_n^2 \varepsilon_{n+3} = 4\nu_n \sigma_{n+3} = 0$$

for $n \geq 9$. Then we can take *extensions* $\overline{\eta_n \circ \varepsilon_{n+1}} \in [\Sigma^{n+7}\mathbb{C}P^2, S^n]$ of $\eta_n \circ \varepsilon_{n+1}$ for $n \geq 9$.

Proposition 8. *(1)* $[\Sigma^{14}\mathbb{C}P^2, S^7] = \mathbb{Z}_8\{\overline{\sigma' \circ \eta_{14}^2 + \eta_7 \circ \varepsilon_8}\} \oplus \mathbb{Z}_2^2\{\overline{\nu}_7 \circ \nu_{15} \circ \Sigma^{14}p, \nu_7^2 \circ g_{13}(\mathbb{C})\}$.
Relation: $2\overline{\sigma' \circ \eta_{14}^2 + \eta_7 \circ \varepsilon_8} = x\zeta_7 \circ \Sigma^{14}p$ *for an odd* x.
(2) $[\Sigma^{15}\mathbb{C}P^2, S^8] = \mathbb{Z}_8\{\overline{(\Sigma\sigma') \circ \eta_{15}^2 + \eta_8 \circ \varepsilon_9}\} \oplus \mathbb{Z}_2^2\{\overline{\nu}_8 \circ \nu_{16} \circ \Sigma^{15}p, \nu_8^2 \circ g_{14}(\mathbb{C})\}$.
Relation: $2\overline{(\Sigma\sigma') \circ \eta_{15}^2 + \eta_8 \circ \varepsilon_9} = x\zeta_8 \circ \Sigma^{15}p$ *for an odd* x.
(3) $[\Sigma^{16}\mathbb{C}P^2, S^9] = \mathbb{Z}_8\{\overline{\eta_9 \circ \varepsilon_{10}}\} \oplus \mathbb{Z}_2^2\{\overline{\nu}_9 \circ \nu_{17} \circ \Sigma^{16}p, \nu_9^2 \circ g_{15}(\mathbb{C})\}$.
Relation: $2\overline{\eta_9 \circ \varepsilon_{10}} = x\zeta_9 \circ \Sigma^{16}p$ *for an odd* x.

Proof. By (3.1) and Lemma 2, we have

$$0 \to \mathbb{Z}_4\{\zeta_7\} \oplus \mathbb{Z}_2\{\overline{\nu}_7 \circ \nu_{15}\} \oplus \mathbb{Z}_7 \oplus \mathbb{Z}_9 \xrightarrow{\Sigma^{14}p^*} [\Sigma^{14}\mathbb{C}P^2, S^7] \xrightarrow{\Sigma^{14}i^*} \mathbb{Z}_2^2\{\nu_7^3, \sigma' \circ \eta_{14}^2 + \eta_7 \circ \varepsilon_8\} \to 0$$

$$0 \to \mathbb{Z}_4\{\zeta_8\} \oplus \mathbb{Z}_2\{\overline{\nu}_8 \circ \nu_{16}\} \oplus \mathbb{Z}_7 \oplus \mathbb{Z}_9 \xrightarrow{\Sigma^{15}p^*} [\Sigma^{15}\mathbb{C}P^2, S^8] \xrightarrow{\Sigma^{15}i^*} \mathbb{Z}_2^2\{\nu_8^3, (\Sigma\sigma') \circ \eta_{15}^2 + \eta_8 \circ \varepsilon_9\} \to 0$$

$$0 \to \mathbb{Z}_4\{\zeta_9\} \oplus \mathbb{Z}_2\{\overline{\nu}_9 \circ \nu_{17}\} \oplus \mathbb{Z}_7 \oplus \mathbb{Z}_9 \xrightarrow{\Sigma^{16}p^*} [\Sigma^{16}\mathbb{C}P^2, S^9] \xrightarrow{\Sigma^{16}i^*} \mathbb{Z}_2^2\{\nu_9^3, \eta_9 \circ \varepsilon_{10}\} \to 0.$$

By [28, Lemma 5.14], we have $\Sigma((\Sigma\sigma') \circ \eta_{15}^2) = (\Sigma^2\sigma') \circ \eta_{16}^2 = (2\sigma_9) \circ \eta_{16}^2 = 0$. Then we obtain

$$\Sigma(\overline{(\Sigma\sigma') \circ \eta_{15}^2 + \eta_8 \circ \varepsilon_9}) \equiv \overline{\eta_9 \circ \varepsilon_{10}} \mod \zeta_9 \circ \Sigma^{16}p, \bar{\nu}_9 \circ \nu_{17} \circ \Sigma^{16}p.$$

This implies that $2\overline{\eta_9 \circ \varepsilon_{10}} = 2\iota_9 \circ \overline{\eta_9 \circ \varepsilon_{10}}$ by [10, Theorem 6.7]. By [28, Proposition 1.9, Lemma 9.1], we obtain

$$2\iota_9 \circ \overline{\eta_9 \circ \varepsilon_{10}} \in \{2\iota_9, \eta_9 \circ \varepsilon_{10}, \eta_{18}\} \circ \Sigma^{16}p \ni \zeta_9 \circ \Sigma^{16}p \mod 2\zeta_9 \circ \Sigma^{16}p,\ \overline{\nu}_9 \circ \nu_{17} \circ \Sigma^{16}p$$

that is,

$$2\overline{\eta_9 \circ \varepsilon_{10}} \equiv x\zeta_9 \circ \Sigma^{16}p \mod \overline{\nu}_9 \circ \nu_{17} \circ \Sigma^{16}p$$

for an odd x. Since $2(\nu_9^2 \circ g_{15}(\mathbb{C})) = 2(\nu_9^2 \circ \Sigma g_{14}(\mathbb{C})) = (2\nu_9^2) \circ g_{15}(\mathbb{C}) = 0$ by [28, (2.1)], $\nu_9^2 \circ g_{15}(\mathbb{C})$ is of order 2. By Lemma 3, we complete the proof. □

By [28, (7.21)], we have $P(\iota_{21}) \circ \eta_{19} = P(\eta_{21}) = 2\sigma_{10} \circ \nu_{17}$. Then we have $2P(\iota_{21}) \circ \eta_{19} = 4\sigma_{10} \circ \nu_{17} = 0$ [28, Theorem 7.3]. Thus we can take an *extension* $P(\iota_{21}) \circ \overline{2\iota_{10}} \in [\Sigma^{17}\mathbb{C}P^2, S^{10}]$ of $2P(\iota_{21})$.

Proposition 9. *(1)* $[\Sigma^{17}\mathbb{C}P^2, S^{10}] = \mathbb{Z}\{P(\iota_{21})\circ\overline{2\iota_{19}}\}\oplus\mathbb{Z}_8\{\overline{\eta_{10}\circ\varepsilon_{11}}\}\oplus\mathbb{Z}_2\{\nu_{10}^2\circ g_{16}(\mathbb{C})\}\oplus\mathbb{Z}_7\oplus\mathbb{Z}_9$.
(2) $[\Sigma^{18}\mathbb{C}P^2, S^{11}] = \mathbb{Z}_8\{\overline{\eta_{11}\circ\varepsilon_{12}}\}\oplus\mathbb{Z}_2\{\nu_{11}^2\circ g_{17}(\mathbb{C})\}\oplus\mathbb{Z}_7\oplus\mathbb{Z}_9$.
(3) $[\Sigma^{19}\mathbb{C}P^2, S^{12}] = \mathbb{Z}\{P(\iota_{25})\circ\Sigma^{19}p\}\oplus\mathbb{Z}_8\{\overline{\eta_{12}\circ\varepsilon_{13}}\}\oplus\mathbb{Z}_2\{\nu_{12}^2\circ g_{18}(\mathbb{C})\}\oplus\mathbb{Z}_7\oplus\mathbb{Z}_9$.
(4) For $n\geq 13$, $[\Sigma^{n+7}\mathbb{C}P^2, S^n] = \mathbb{Z}_8\{\overline{\eta_n\circ\varepsilon_{n+1}}\}\oplus\mathbb{Z}_2\{\nu_n^2\circ g_{n+6}(\mathbb{C})\}\oplus\mathbb{Z}_7\oplus\mathbb{Z}_9$.

Proof. (1) By (3.1) and Lemma 2, we have

$$0\to\mathbb{Z}_4\{\zeta_{10}\}\oplus\mathbb{Z}_7\oplus\mathbb{Z}_9\xrightarrow{\Sigma^{17}p^*}[\Sigma^{17}\mathbb{C}P^2, S^{10}]\xrightarrow{\Sigma^{17}i^*}\mathbb{Z}\{2P(\iota_{21})\}\oplus\mathbb{Z}_2^2\{\nu_{10}^3,\eta_{10}\circ\varepsilon_{11}\}\to 0.$$

By proposition 8, we have

$$2\overline{\eta_{10}\circ\varepsilon_{11}}\equiv x\zeta_{10}\circ\Sigma^{17}p \mod \bar{\nu}_{10}\circ\nu_{18}\circ\Sigma^{17}p$$

for an odd x. Since ν_{11}^2 is of order 2, we have $\nu_{11}^2\circ g_{17}(\mathbb{C})$ is of order 2. Thus we have

$$[\Sigma^{17}\mathbb{C}P^2, S^{10}] = \mathbb{Z}\{\overline{2P(\iota_{21})}\}\oplus\mathbb{Z}_8\{\overline{\eta_{10}\circ\varepsilon_{11}}\}\oplus\mathbb{Z}_2\{\nu_{10}^2\circ g_{16}(\mathbb{C})\}\oplus\mathbb{Z}_7\oplus\mathbb{Z}_9.$$

(2) By (3.1) and Lemma 2, we have

$$0\to\mathbb{Z}_4\{\zeta_{11}\}\oplus\mathbb{Z}_7\oplus\mathbb{Z}_9\xrightarrow{\Sigma^{18}p^*}[\Sigma^{18}\mathbb{C}P^2, S^{11}]\xrightarrow{\Sigma^{18}i^*}\mathbb{Z}_2^2\{\nu_{11}^3,\eta_{11}\circ\varepsilon_{12}\}\to 0.$$

By (1), we obtain

$$[\Sigma^{18}\mathbb{C}P^2, S^{11}] = \mathbb{Z}_8\{\overline{\eta_{11}\circ\varepsilon_{12}}\}\oplus\mathbb{Z}_2\{\nu_{11}^2\circ g_{17}(\mathbb{C})\}\oplus\mathbb{Z}_7\oplus\mathbb{Z}_9.$$

Similarly, we can determine group structures of $[\Sigma^{n+7}\mathbb{C}P^2, S^n]$ for $n\geq 12$. This completes the proof. □

For odd prime p, we have an isomorphism

$$[\Sigma^{n+7}\mathbb{C}P^2, S^n]_{(p)}\cong\pi_{n+9}(S^n)_{(p)}\oplus\pi_{n+11}(S^n)_{(p)} \tag{2}$$

by (2.2). From [28, Chapter 13], then, we have the following:

Proposition 10. *The odd primary components of* $[\Sigma^{n+7}\mathbb{C}P^2, S^n]$ *are*
(1) $[\Sigma^{9}\mathbb{C}P^2, S^2] = \mathbb{Z}_3\{\eta_2\circ\alpha_3(3)\circ\Sigma^9p\}$
(2) $[\Sigma^{10}\mathbb{C}P^2, S^3] = \mathbb{Z}_3\{\alpha_3(3)\circ\Sigma^{10}p\}\oplus\mathbb{Z}_7\{\alpha_{1,7}(3)\circ\Sigma^{10}p\}$.
(3) $[\Sigma^{11}\mathbb{C}P^2, S^4] = \mathbb{Z}_3\{\alpha_3(4)\circ\Sigma^{11}p\}\oplus\mathbb{Z}_7\{\alpha_{1,7}(4)\circ\Sigma^{11}p\}$.
(4) $[\Sigma^{n+7}\mathbb{C}P^2, S^n] = \mathbb{Z}_7\{\alpha_{1,7}(n)\circ\Sigma^{n+7}p\}\oplus\mathbb{Z}_9\{\alpha_3'(n)\}$ *for* $n\geq 5$.

5. Determination of $[\Sigma^{n+8}\mathbb{C}P^2, S^n]$

In this section, we determine the generators of the 2-primary components of n-th cohomotopy groups of $(n+8)$-fold suspended complex projective planes. We analyze the short exact sequence

$$0\to Coker\,\eta_{n+11}^*\xrightarrow{\Sigma^n p^*}[\Sigma^{n+8}\mathbb{C}P^2, S^n]\xrightarrow{\Sigma^n i^*}Ker\,\eta_{n+10}^*\to 0 \tag{3.1}$$

induced by (1.1).

$n=$	2	3	4	5	6	7, 8, 9	10	11	12	13	$14 \geq$
$Coker\,\eta^*_{n+11}$	$\mathbb{Z}_2^2$	0	$\mathbb{Z}_2$	0	$\mathbb{Z}_8$	0	$\mathbb{Z}_4$	$\mathbb{Z}_2$	$\mathbb{Z}_2$	$\mathbb{Z}_2$	0
generators	$\eta_2\mu'$ $\eta_2\nu'\bar{\nu}_6$	0	$\nu_4\mu_7$	0	$P(\sigma_{13})$	0	$P(\nu_{21})$	θ'	θ	$\Sigma\theta$	0

Lemma 4. *(1) For the homomorphisms $\eta^*_{n+11} : \pi^n_{n+11} \to \pi^n_{n+12}$, we have the following table of the cokernel of η^*_{n+11} :*

(2) For the homomorphisms $\eta^*_{n+10} : \pi^n_{n+10} \to \pi^n_{n+11}$, we have the following table of the kernel of η^*_{n+10} :

$n=$	2	3	4	5, 6	7	8	9	10	11	$12 \geq$
$Ker\,\eta^*_{n+10}$	0	$\mathbb{Z}_2$	$\mathbb{Z}_4 \oplus \mathbb{Z}_2$	$\mathbb{Z}_4$	$\mathbb{Z}_8$	$\mathbb{Z}_8^2$	$\mathbb{Z}_8$	$\mathbb{Z}_4$	$\mathbb{Z}_2$	0
generators	0	$\eta_3^2\varepsilon_5$	$2(\nu_4\sigma')$ $\eta_4^2\varepsilon_6$	$2\nu_n\sigma_{n+3}$	$\nu_7\sigma_{10}$	$\sigma_8\nu_{15}$ $\nu_8\sigma_{11}$	$\sigma_9\nu_{16}$	$\sigma_{10}\nu_{17}$	$\sigma_{11}\nu_{18}$	0

By Lemma 4, we have the following

Proposition 11. $[\Sigma^{10}\mathbb{C}P^2, S^2] = \mathbb{Z}_2^2\{\eta_2 \circ \mu' \circ \Sigma^{10}p, \eta_2 \circ \nu' \circ \bar{\nu}_6 \circ \Sigma^{10}p\} \oplus \mathbb{Z}_{21}$.

By [28, (7.12)], we have $\varepsilon' \circ \eta_{13} = \nu' \circ \varepsilon_6$ and $\nu' \circ \varepsilon_6$ is of order 2. Then we can take an extension $\overline{\eta_3^2 \circ \varepsilon_5} \in [\Sigma^{11}\mathbb{C}P^2, S^3]$ of $2\varepsilon' = \eta_3^2 \circ \varepsilon_5$ [28, Lemma 6.6]. Similarly, we can take an extension $\overline{\eta_4^2 \circ \varepsilon_6} \in [\Sigma^{12}\mathbb{C}P^2, S^4]$ of $\eta_4^2 \circ \varepsilon_6$. Then we obtain the following.

Proposition 12. $[\Sigma^{11}\mathbb{C}P^2, S^3] = \mathbb{Z}_2\{\overline{\eta_3^2 \circ \varepsilon_5}\} \oplus \mathbb{Z}_3$.

Proposition 13. $[\Sigma^{12}\mathbb{C}P^2, S^4] = \mathbb{Z}_8\{\nu_4 \circ \sigma' \circ \overline{2\iota_{14}}\} \oplus \mathbb{Z}_2\{\overline{\eta_4^2 \circ \varepsilon_6}\}$.

Proof. By Lemma 4, we have a short exact sequence

$$0 \to \mathbb{Z}_2\{\nu_4 \circ \mu_7\} \xrightarrow{\Sigma^{12}p^*} [\Sigma^{12}\mathbb{C}P^2, S^4] \xrightarrow{\Sigma^{12}i^*} \mathbb{Z}_4\{2(\nu_4 \circ \sigma')\} \oplus \mathbb{Z}_2\{2\Sigma\varepsilon'\} \to 0.$$

By [11, Proposition 3.7], we obtain

$$4\nu_4 \circ \sigma' \circ \overline{2\iota_{14}} = \nu_4 \circ \mu_7 \circ \Sigma^{12}p.$$

This complete the proof. □

By [28, p.152], $\nu_5 \circ \sigma_8 \circ \eta_{15} = \nu_5 \circ \varepsilon_8$ is of order 2. Then we can take an extension $\nu_5 \circ \sigma_8 \circ \overline{2\iota_{15}} \in [\Sigma^{13}\mathbb{C}P^2, S^5]$ of $2\nu_5 \circ \sigma_8$. Similarly, we can take an extension $\nu_6 \circ \sigma_9 \circ \overline{2\iota_{16}} \in [\Sigma^{14}\mathbb{C}P^2, S^6]$ of $2\nu_6 \circ \sigma_9$. By Lemma 4, we have the following

Proposition 14. *(1)* $[\Sigma^{13}\mathbb{C}P^2, S^5] = \mathbb{Z}_4\{\nu_5 \circ \sigma_8 \circ \overline{2\iota_{15}}\} \oplus \mathbb{Z}_9$.
(2) $[\Sigma^{14}\mathbb{C}P^2, S^6] = \mathbb{Z}_8\{P(\sigma_{13}) \circ \Sigma^{14}p\} \oplus \mathbb{Z}_4\{\nu_5 \circ \sigma_8 \circ \overline{2\iota_{15}}\} \oplus \mathbb{Z}_9 \oplus \mathbb{Z}_3$.

Proof. By Lemma 4, we have (1). So we only prove (2). Consider the following commutative diagram induced by Lemma 4.

$$\begin{tikzcd}
0 \arrow[r] & 0 \arrow[r] \arrow[d] & {[\Sigma^{13}\mathbb{C}P^2, S^5]} \arrow[r, "\Sigma^{13}i^*"] \arrow[d, "\Sigma_1"] & \mathbb{Z}_4\{2\nu_5 \circ \sigma_8\} \arrow[r] \arrow[d, "\Sigma_2"] & 0 \\
0 \arrow[r] & \mathbb{Z}_8\{P(\sigma_{13})\} \arrow[r, "\Sigma^{13}p^*"] & {[\Sigma^{14}\mathbb{C}P^2, S^6]} \arrow[r, "\Sigma^{14}i^*"] & \mathbb{Z}_4\{2\nu_6 \circ \sigma_9\} \arrow[r] & 0
\end{tikzcd}$$

Since $\Sigma^{13}i^*$ and Σ_2 are isomorphic, the second row is split. This complete the proof. □

By [28, (5.9)], $\nu_n \circ \sigma_{n+3} \circ \eta_{n+10} = 0$ for $n \geq 7$. Then we can take an extension $\overline{\nu_n \circ \sigma_{n+3}} \in [\Sigma^{n+8}\mathbb{C}P^2, S^n]$ of $\nu_n \circ \sigma_{n+3}$. Similarly, we have $\sigma_m \circ \nu_{m+7} \circ \eta_{m+10} = \nu_m \circ \sigma_{m+3} \circ \eta_{m+10} = 0$ for $m \geq 8$. Then we can take extensions $\overline{\sigma_m \circ \nu_{m+7}}, \overline{\nu_m \circ \sigma_{m+3}} \in [\Sigma^{m+8}\mathbb{C}P^2, S^m]$ of $\sigma_m \circ \nu_{m+7}$ and $\nu_m \circ \sigma_{m+3}$ respectively. By Lemma 4, we have the following.

Proposition 15. *(1)* $[\Sigma^{15}\mathbb{C}P^2, S^7] = \mathbb{Z}_8\{\overline{\nu_7 \circ \sigma_{10}}\} \oplus \mathbb{Z}_3$.
(2) $[\Sigma^{16}\mathbb{C}P^2, S^8] = \mathbb{Z}_8^2\{\overline{\sigma_8 \circ \nu_{15}}, \overline{\nu_8 \circ \sigma_{11}}\} \oplus \mathbb{Z}_3^2$.
(3) $[\Sigma^{17}\mathbb{C}P^2, S^9] = \mathbb{Z}_8\{\overline{\sigma_9 \circ \nu_{16}}\} \oplus \mathbb{Z}_3$.

Lemma 5. $\{2\sigma_{10} \circ \nu_{17}, 2\iota_{20}, \eta_{20}\}_1 = 2P(\nu_{21})$.

Proof. We recall that $\pi_{22}^{10} = \mathbb{Z}_4\{P(\nu_{21})\}$ [28, Proposition 7.6]. By [28, Proposition 2.6, (7.21)], we have

$$H\{2\sigma_{10} \circ \nu_{17}, 2\iota_{20}, \eta_{20}\}_1 = \eta_{19}^3 = 4\nu_{19}.$$

Since $(H \circ P)(\nu_{21}) = \pm 2\nu_{19}$ [28, Proposition 2.7], we obtain the result. □

Proposition 16. $[\Sigma^{18}\mathbb{C}P^2, S^{10}] = \mathbb{Z}_8\{\sigma_{10} \circ g_{17}(\mathbb{C})\} \oplus \mathbb{Z}_2\{2\sigma_{10} \circ g_{17}(\mathbb{C}) - P(\nu_{21}) \circ \Sigma^{18}p\}$.

Proof. By Lemma 4, we have a short exact sequence

$$0 \to \mathbb{Z}_4\{P(\nu_{21})\} \xrightarrow{\Sigma^{18}p^*} [\Sigma^{18}\mathbb{C}P^2, S^{10}] \xrightarrow{\Sigma^{18}i^*} \mathbb{Z}_4\{\sigma_{10} \circ \nu_{17}\} \to 0.$$

By [28, Proposition 1.9], we have

$$\begin{aligned} 4\sigma_{10} \circ g_{17}(\mathbb{C}) &= 2\sigma_{10} \circ \nu_{17} \circ \overline{2\iota_{20}} \\ &\in \{2\sigma_{10} \circ \nu_{17}, 2\iota_{20}, \eta_{20}\}_1 \circ \Sigma^{18}p \\ &= 2P(\nu_{21}) \circ \Sigma^{18}p. \end{aligned}$$

This completes the proof. □

Proposition 17. $[\Sigma^{19}\mathbb{C}P^2, S^{11}] = \mathbb{Z}_2^2\{\sigma_{11} \circ g_{18}(\mathbb{C}), \theta' \circ \Sigma^{19}p\}$.

Proof. By Lemma 4, we have a short exact sequence

$$0 \to \mathbb{Z}_2\{\theta'\} \xrightarrow{\Sigma^{19}p^*} [\Sigma^{19}\mathbb{C}P^2, S^{11}] \xrightarrow{\Sigma^{19}i^*} \mathbb{Z}_2\{\sigma_{11} \circ \nu_{18}\} \to 0.$$

We consider 2-primary EHP-sequence

$$[\Sigma^{20}\mathbb{C}P^2, S^{21}] \xrightarrow{P} [\Sigma^{18}\mathbb{C}P^2, S^{10}] \xrightarrow{\Sigma} [\Sigma^{19}\mathbb{C}P^2, S^{11}] \xrightarrow{H}$$

$$\xrightarrow{H} [\Sigma^{19}\mathbb{C}P^2, S^{21}] \xrightarrow{P} [\Sigma^{17}\mathbb{C}P^2, S^{10}]$$

where $[\Sigma^{20}\mathbb{C}P^2, S^{21}] = \mathbb{Z}_4\{\nu_{21} \circ \Sigma^{20}p\}$ and $[\Sigma^{19}\mathbb{C}P^2, S^{21}] = \mathbb{Z}\{\overline{2\iota_{21}}\}$ [11, p.110, Proposition 3.1]. Then $P(\nu_{21} \circ \Sigma^{20}p) = P(\nu_{21}) \circ \Sigma^{18}p$ and $P(\overline{2\iota_{21}}) \equiv P(\iota_{21}) \circ \overline{2\iota_{19}} \mod \nu_{10}^2 \circ g_{16}(\mathbb{C}), 2\overline{\eta_{10} \circ \varepsilon_{11}}$. So, two P-homomorphisms are monomorphisms. Thus $[\Sigma^{19}\mathbb{C}P^2, S^{11}] \cong \mathbb{Z}_2^2$. This completes the proof. □

By Lemma 4, we have the following.

Proposition 18. *(1)* $[\Sigma^{20}\mathbb{C}P^2, S^{12}] = \mathbb{Z}_2\{\theta \circ \Sigma^{20}p\} \oplus \mathbb{Z}_3$.
(2) $[\Sigma^{21}\mathbb{C}P^2, S^{13}] = \mathbb{Z}_2\{(\Sigma\theta) \circ \Sigma^{21}p\} \oplus \mathbb{Z}_3$.
(3) $[\Sigma^{n+8}\mathbb{C}P^2, S^n] = \mathbb{Z}_3$ *for* $n \geq 14$.

For odd prime p, we have an isomorphism

$$[\Sigma^{n+8}\mathbb{C}P^2, S^n]_{(p)} \cong \pi_{n+10}(S^n)_{(p)} \oplus \pi_{n+12}(S^n)_{(p)} \tag{3}$$

by (2.2). From [28, Chapter 13], then, we have the following:

Proposition 19. *The odd primary components of* $[\Sigma^{n+8}\mathbb{C}P^2, S^n]$ *are*
(1) $[\Sigma^{10}\mathbb{C}P^2, S^2] = \mathbb{Z}_3\{\eta_2 \circ \alpha_3(3) \circ \Sigma^{10}p\} \oplus \mathbb{Z}_5\{\eta_2 \circ \alpha_{1,7}(3) \circ \Sigma^{10}p\}$.
(2) $[\Sigma^{11}\mathbb{C}P^2, S^3] = \mathbb{Z}_3\{\alpha_1(3) \circ \overline{\alpha_2(6)}\}$.
(3) $[\Sigma^{12}\mathbb{C}P^2, S^4] = \mathbb{Z}_3^2\{\alpha_1(3) \circ \overline{\alpha_2(6)}, [\iota_4, \iota_4] \circ \overline{\alpha_2(7)}\}$.
(4) $[\Sigma^{13}\mathbb{C}P^2, S^5] = \mathbb{Z}_9\{\overline{\beta_1(5)}\}$.
(5) $[\Sigma^{14}\mathbb{C}P^2, S^6] = \mathbb{Z}_9\{\overline{\beta_1(6)}\} \oplus \mathbb{Z}_3\{[\iota_6, \iota_6] \circ \alpha_2(11) \circ \Sigma^{14}p\} \oplus \mathbb{Z}_5\{[\iota_6, \iota_6] \circ \alpha_{1,5}(11) \circ \Sigma^{14}p\}$.
(6) $[\Sigma^{15}\mathbb{C}P^2, S^7] = \mathbb{Z}_3\{\overline{\beta_1(7)}\}$.
(7) $[\Sigma^{16}\mathbb{C}P^2, S^8] = \mathbb{Z}_3^2\{\overline{\beta_1(8)}, [\iota_8, \iota_8] \circ \overline{\alpha_1(15)}\}$.
(8) $[\Sigma^{17}\mathbb{C}P^2, S^9] = \mathbb{Z}_3\{\overline{\beta_1(9)}\}$.
(9) $[\Sigma^{18}\mathbb{C}P^2, S^{10}] = \mathbb{Z}_3^2\{\overline{\beta_1(9)}, [\iota_{10}, \iota_{10}] \circ \alpha_1(19) \circ \Sigma^{18}p\}$.
(10) $[\Sigma^{n+8}\mathbb{C}P^2, S^n] = \mathbb{Z}_3\{\overline{\beta_1(n)}\}$ *for* $n \geq 11$.

6. Homotopy groups of $Map_*(\mathbb{C}P^2, \mathbb{C}P^2, *)$

The complex projective plane $\mathbb{C}P^2$ is a base space of a S^1-bundle $S^1 \to S^5 \xrightarrow{p} \mathbb{C}P^2$. So, we have an isomorphism

$$p_* : [\Sigma^n\mathbb{C}P^2, S^5] \to [\Sigma^n\mathbb{C}P^2, \mathbb{C}P^2].$$

From the adjointness [15], we obtain an isomorphism

$$\pi_n(map_*(\mathbb{C}P^2, \mathbb{C}P^2; *)) \cong [\Sigma^n\mathbb{C}P^2, \mathbb{C}P^2].$$

McGibbon [20] showed that homotopy class $[\mathbb{C}P^n, \mathbb{C}P^n] \cong \mathbb{Z}$ which is determined by using homomorphism between homology groups. We denote the n-th homotopy group $\pi_n(Map_*(\mathbb{C}P^2, \mathbb{C}P^2; *))$ by $\pi_n(map_*(\mathbb{C}P^2, \mathbb{C}P^2))$. By the results of section 3, 4, 5 and [11], we obtain the following:

Theorem 1. $\pi_4(map_*(\mathbb{C}P^2, \mathbb{C}P^2)) \cong \mathbb{Z}_4\{\nu_5 \circ \Sigma^4 p\} \oplus \mathbb{Z}_3\{\alpha_1(5) \circ \Sigma^4 p\}$.
$\pi_5(map_*(\mathbb{C}P^2, \mathbb{C}P^2)) = 0$.
$\pi_6(map_*(\mathbb{C}P^2, \mathbb{C}P^2)) \cong \mathbb{Z}_4\{\nu_5 \circ \overline{2\iota_8}\} \oplus \mathbb{Z}_3\{\overline{\alpha_1(5)}\}$.
$\pi_7(map_*(\mathbb{C}P^2, \mathbb{C}P^2)) \cong \mathbb{Z}_2\{\nu_5^2 \circ \Sigma^7 p\}$.
$\pi_8(map_*(\mathbb{C}P^2, \mathbb{C}P^2)) \cong \mathbb{Z}_4\{\overline{\nu_5 \circ \eta_8^2}\} \oplus \mathbb{Z}_3\{\alpha_2(5) \circ \Sigma^8 p\} \oplus \mathbb{Z}_5\{\alpha_1'(5) \circ \Sigma^8 p\}$.
$\pi_9(map_*(\mathbb{C}P^2, \mathbb{C}P^2)) \cong \mathbb{Z}_4\{\nu_5 \circ g_8(\mathbb{C})\}$.
$\pi_{10}(map_*(\mathbb{C}P^2, \mathbb{C}P^2)) \cong \mathbb{Z}_2\{\nu_5^3 \circ \Sigma^{10} p\} \oplus \mathbb{Z}_4\{\overline{\sigma'''}\} \oplus \mathbb{Z}_3\{\overline{\alpha_2(5)}\} \oplus \mathbb{Z}_5\{\overline{\alpha_1'(5)}\}$.
$\pi_{11}(map_*(\mathbb{C}P^2, \mathbb{C}P^2)) \cong \mathbb{Z}_4\{\nu_5 \circ \sigma_8 \circ \Sigma^{11} p\} \oplus \mathbb{Z}_9\{\beta_1(5) \circ \Sigma^{11} p\}$.
$\pi_{12}(map_*(\mathbb{C}P^2, \mathbb{C}P^2)) \cong \mathbb{Z}_4\{\zeta_5 \circ \Sigma^{12} p\} \oplus \mathbb{Z}_2^2\{\nu_5 \circ \overline{\nu}_8 \circ \Sigma^{12} p, \nu_5^2 \circ g_{11}(\mathbb{C})\} \oplus \mathbb{Z}_9\{\alpha_3(5) \circ \Sigma^{12} p\} \oplus \mathbb{Z}_7$.
$\pi_{13}(map_*(\mathbb{C}P^2, \mathbb{C}P^2)) \cong \mathbb{Z}_4\{\nu_5 \circ \sigma_8 \circ \overline{2\iota_{15}}\} \oplus \mathbb{Z}_9\{\overline{\beta_1(5)}\}$.

7. Applications : Classifying path-components of mapping spaces and Cyclic maps

In this section, we apply the results obtained in Sections 3, 4 and 5 to classify homotopy types of path-components of certain mapping spaces and compute certain generalized Gottlieb groups.

The term fibration is used for a Hurewicz fibration, that is a map with the homotopy lifting property with respect to all spaces [19, p.49]. It is well known that the evaluation map $w_f : map(X, Y; f) \to Y$, $w_f(g) = g(*)$, is a fibration [7, Lemma 8.15]. For fibrations $p : E_1 \to B$ and $q : E_2 \to B$, p and q are said to be *fiber homotopy equivalent* if there is a homotopy equivalence $h : E_1 \to E_2$ such that $q \circ h = p$ [19, p.52].

Here we remind several results of the generalized Whitehead product [1]. If α and β are homotopy classes, then the Whitehead product of α and β is denoted by $[\alpha, \beta]$. A map $f : X \to Y$ is cyclic if there is a map $F : X \times Y \to Y$, called affiliated map, such that $F(x, *) = f(x)$ and $F(*, y) = y$. We denote the set of cyclic map from X to Y by $G(X, Y)$ and it has group structure if X is a co-H-group [29].

We recall the following equivalent statements due to [8, Lemma 2] and [16, Theorem 3.7]:

Theorem 1. *Let ΣX and ΣY be CW complexes with non-degenerate basepoints and ΣX is finite CW complex. Then the following are equivalent.*
(A) the map $f : \Sigma X \to \Sigma Y$ is cyclic
(B) $[f, id_{\Sigma Y}] = 0$ where $[\ ,\ ]$ is the generalized Whitehead product
(C) the evaluation fibration $w_f : map(\Sigma X, \Sigma Y; f) \to \Sigma Y$ has a section

(D) the evaluation fibration $w_f : map(\Sigma X, \Sigma Y; f) \to \Sigma Y$ *is fibre-homotopy equivalent to* $w_* : Map(\Sigma X, \Sigma Y; *) \to Y$

Here is a connection of path components of mapping spaces and cyclic maps.

Theorem 2. [17, Theorem 3.10] *Suppose* X *is a CW co-H-space and* Y *is any CW complex. Let* $d \in G(X, Y)$ *be any cyclic map. Then for each map* $f : X \to Y$*, we have* $map(X, Y; f) \simeq map(X, Y; f + d)$*. If* X *is a finite co-H-space then the corresponding evaluation fibrations* w_f *and* w_{f+d} *are fibre-homotopy equivalent.*

The following Theorem shows a relation between the generaized Whitehead product and evlauation fibration [8, Theorem 1]

Theorem 3. *Given a pair of homotopy classes* $\alpha = [f], \beta = [g] \in [\Sigma A, \Sigma B]$ *such that at least one of the identities* $[\alpha, \iota_{\Sigma B}] = \pm[\beta, \iota_{\Sigma B}]$ *holds. Then the evaluation fibrations* $w_f : map(\Sigma A, \Sigma B; f) \to \Sigma B)$ *and* $w_g : map(\Sigma A, \Sigma B; g) \to \Sigma B)$ *are fibre homotopy equivalent.*

We recall Proposition 4.4 of [11].

Proposition 20. *(1)* $[\Sigma\mathbb{C}P^2, S^2] \cong \mathbb{Z}\{\eta_2 \circ \overline{2\iota_3}\}$*.*
(2) $[\Sigma^2\mathbb{C}P^2, S^3] \cong \mathbb{Z}_2\{\nu' \circ \Sigma^2 p\} \oplus \mathbb{Z}_3\{\alpha_1(3) \circ \Sigma^2 p\}$*.*
(3) $[\Sigma^3\mathbb{C}P^2, S^4] \cong \mathbb{Z}\{\nu_4 \circ \Sigma^3 p\} \oplus \mathbb{Z}_2\{\Sigma\nu' \circ \Sigma^3 p\} \oplus \mathbb{Z}_3\{\alpha_1(4) \circ \Sigma^3 p\}$*.*
(4) $[\Sigma^n\mathbb{C}P^2, S^{n+1}] \cong \mathbb{Z}_4\{\nu_{n+1} \circ \Sigma^n p\} \oplus \mathbb{Z}_3\{\alpha_1(n+1) \circ \Sigma^n p\}$ *for* $n \geq 4$*.*

Let G be an abelian group and let S be a subset of G. We denote the smallest subgroup of G containing a subset $S \subset G$ by $\langle S \rangle$. We denote $\omega_f : map(X, Y; f) \to Y$ by ω_f for $f : X \to Y$.

Theorem 4. *(1) For each* $[f], [g] \in [\Sigma\mathbb{C}P^2, S^2]$*, the evaluation fibrations* w_f *and* w_g *are fibre homotopy equivalent.*
(2) For each $[f], [g] \in [\Sigma^2\mathbb{C}P^2, S^3]$*, the evaluation fibrations* w_f *and* w_g *are fibre homotopy equivalent.*
(3) (a) For each $[f], [g] \in \left\langle 2\nu_4 \circ \Sigma^3 p, \Sigma\nu' \circ \Sigma^3 p \right\rangle$*, the evaluation fibrations* w_f *and* w_g *are fibre homotopy equivalent.*
(b) For each $[f] \in \left\langle 2\nu_4 \circ \Sigma^3 p, \Sigma\nu' \circ \Sigma^3 p \right\rangle$ *and* $[g] \in \{(2n+1)\nu_4 \circ \Sigma^3 p \,|\, n \in \mathbb{Z}\}$*, the evaluation fibrations* w_g *and* w_{f+g} *are fibre homotopy equivalent.*
(c) For each $[f] \in \left\langle 2\nu_4 \circ \Sigma^3 p, \Sigma\nu' \circ \Sigma^3 p \right\rangle$ *and* $[h] \in \{\alpha_1(4) \circ \Sigma^3 p, 2\alpha_1(4) \circ \Sigma^3 p\}$*, the evaluation fibrations* w_h *and* w_{h+f} *are fibre homotopy equivalent.*
(4) For each $[f], [g] \in [\Sigma^4\mathbb{C}P^2, S^5]$*, the evaluation fibrations* w_f *and* w_g *are fibre homotopy equivalent.*
(5) (a) The evaluation fibrations w_* *and* w_f *are fiber homotopy equivalent where* $[f] = 2\nu_6 \circ \Sigma^5 p$*.*
(b) For each $[f] \in \{\nu_6 \circ \Sigma^5 p, 3\nu_6 \circ \Sigma^5 p\}$ *and* $[g] \in \left\langle 2\nu_6 \circ \Sigma^5 p \right\rangle$*, the evaluation fibrations* w_f *and* w_{f+g} *are fibre homotopy equivalent.*
(c) For each $[f] \in \{\alpha_1(6) \circ \Sigma^5 p, 2\alpha_1(6) \circ \Sigma^5 p\}$ *and* $[g] \in \left\langle 2\nu_6 \circ \Sigma^5 p \right\rangle$*, the evaluation fibrations* w_f *and* w_{f+g} *are fibre homotopy equivalent.*

(6) For each $[f], [g] \in [\Sigma^6\mathbb{C}P^2, S^7]$*, the evaluation fibrations* w_f *and* w_g *are fibre homotopy equivalent.*
(7) (a) Two evaluation fibrations w_f *and* w_g *are fibre homotopy equivalent where* $[f] = \nu_8 \circ \Sigma^7 p$ *and* $[g] = 3\nu_8 \circ \Sigma^7 p$*.*
(b) Two evaluation fibrations w_f *and* w_g *are fibre homotopy equivalent where* $[f] = \alpha_1(9) \circ \Sigma^7 p$ *and* $[g] = 2\alpha_1(9) \circ \Sigma^7 p$*.*
(8) (a) For each $[f], [g] \in \langle \alpha_1(9) \circ \Sigma^8 p, 2\nu_9 \circ \Sigma^8 p \rangle$*, the evaluation fibrations* w_f *and* w_g *are fibre homotopy equivalent.*
(b) For each $[f] \in \{\nu_9 \circ \Sigma^8 p, 3\nu_9 \circ \Sigma^8 p\}$ *and* $[g] \in \langle \alpha_1(9) \circ \Sigma^8 p, 2\nu_9 \circ \Sigma^8 p \rangle$*, the evaluation fibrations* w_f *and* w_{f+g} *are fibre homotopy equivalent.*

Proof. (1) By [13, (13)] and [2, (2.9)], we have $[\iota_2, \eta_2 \circ \overline{2\iota_3}] = 0$. Then we obtain (1) by Theorem 1.
(2) Since S^3 is an H-space, $[\iota_3, f] = 0$ for any $f \in [\Sigma^2\mathbb{C}P^2, S^3]$. Then we obtain (2) by Theorem 1.
(3) By [28, (5.13)] and [11, Proposition 3.4], we have

$$\begin{aligned}
[\iota_4, \nu_4 \circ \Sigma^3 p] &= [\iota_4, \nu_4] \circ \Sigma(\iota_3 \wedge \Sigma^2 p) = 2\nu_4^2 \circ \Sigma^6 p \\
[\iota_4, \Sigma\nu' \circ \Sigma^3 p] &= [\iota_4, \Sigma\nu'] \circ \Sigma(\iota_3 \wedge \Sigma^2 p) = 4\nu_4^2 \circ \Sigma^6 p = 0.
\end{aligned}$$

Then we obtain (a) and (b) by Theorem 1 and 2. By [11, Proposition 3.4]

$$[\iota_4, \alpha_1(4) \circ \Sigma^3 p] = [\iota_4, \alpha_1(4)] \circ \Sigma(\iota_3 \wedge \Sigma^2 p) = [\iota_4, \iota_4] \circ \alpha_1(7) \circ \Sigma^6 p$$

is of order 3. Then we have (c) by Theorem 2 and 3.
(4) By [28, P.48], we have

$$[\iota_5, \nu_5 \circ \Sigma^4 p] = [\iota_5, \nu_5] \circ \Sigma(\iota_4 \wedge \Sigma^3 p) = 0$$

Since $G_8(S^5) = \pi_8(S^5)$ [4, p.428], we obtain

$$[\iota_5, \alpha_1(5) \circ \Sigma^4 p] = [\iota_5, \alpha_1(5)] \circ \Sigma(\iota_4 \wedge \Sigma^3 p) = 0.$$

Thus we have (4) by Theorem 2 and 3.
(5) By [28, p.63], we have

$$[\iota_6, \nu_6 \circ \Sigma^5 p] = [\iota_6, \nu_6] \circ \Sigma(\iota_5 \wedge \Sigma^4 p) = 2\overline{\nu}_6 \circ \Sigma^{10} p.$$

Since $\overline{\nu}_6 \circ \Sigma^{10} p$ is of order 4 [11, Proposition 3.7], we have (a) and (b) by Theorem 1 and 2. By [11, Proposition 3.7],

$$[\iota_6, \alpha_1(6) \circ \Sigma^5 p] = [\iota_6, \alpha_1(6)] \circ \Sigma(\iota_5 \wedge \Sigma^4 p) = [\iota_6, \iota_6] \circ \alpha_1(11) \circ \Sigma^{10} p$$

is of order 3. Then we have (c) by Theorem 2 and 3.
(6) Since S^7 is an H-space, $[\iota_7, f] = 0$ for any $f \in [\Sigma^8\mathbb{C}P^2, S^7]$. Then we obtain (6) by Theorem 1.
(7) By [28, (7.19)], we have

$$[\iota_8, \nu_8 \circ \Sigma^7 p] = [\iota_8, \nu_8] \circ \Sigma(\iota_7 \wedge \Sigma^6 p) = 2\sigma_8 \circ \nu_{15} \circ \Sigma^{14} p - x\nu_8 \circ \sigma_{11} \circ \Sigma^{14} p$$

for an odd x. By Theorem 1 and 2, we have (a) and (b). By Proposition 2,

$$[\iota_8, \alpha_1(8)\circ\Sigma^7 p] = [\iota_8, \alpha_1(8)]\circ\Sigma(\iota_7\wedge\Sigma^6 p) = [\iota_8,\iota_8]\circ\alpha_1(15)\circ\Sigma^{14}p$$

is of order 3. By Theorem 2 and 3, we have (c).
(8) By [28, (7.22)], we have

$$[\iota_9, \nu_9\circ\Sigma^8 p] = [\iota_9,\nu_9]\circ\Sigma(\iota_8\wedge\Sigma^7 p) = \overline{\nu}_9\circ\nu_{17}\circ\Sigma^{16}p.$$

Since $\overline{\nu}_9\circ\nu_{17}\circ\Sigma^{16}p$ is of order 2, we have $[\iota_9, 2\nu_9\circ\Sigma^8 p] = 0$. Since $G_{12}(S^9) = \pi_{12}(S^9)$ [4, p.428], we obtain

$$[\iota_9, \alpha_1(9)\circ\Sigma^8 p] = [\iota_9, \alpha_1(9)]\circ\Sigma(\iota_8\wedge\Sigma^7 p) = 0.$$

By Theorem 2 and 3, we have (b).

□

Corollary 1. *(1)* $G_1(\mathbb{C}P^2, S^2) = Z\{\eta_2\circ\overline{2\iota_3}\}$.
(2) $G_2(\mathbb{C}P^2, S^3) = \mathbb{Z}_2\{\nu'\circ\Sigma^2 p\}\oplus\mathbb{Z}_3\{\alpha_1(3)\circ\Sigma^2 p\}$.
(3) $G_3(\mathbb{C}P^2, S^4) = \mathbb{Z}\{2\nu_4\circ\Sigma^3 p\}\oplus\mathbb{Z}_2\{\Sigma\nu'\circ\Sigma^3 p\}$.
(4) $G_4(\mathbb{C}P^2, S^5) = \mathbb{Z}_4\{\nu_5\circ\Sigma^4 p\}\oplus\mathbb{Z}_3\{\alpha_1(5)\circ\Sigma^4 p\}$.
(5) $G_5(\mathbb{C}P^2, S^6) = \mathbb{Z}_2\{2\nu_6\circ\Sigma^5 p\}$.
(6) $G_6(\mathbb{C}P^2, S^7) = \mathbb{Z}_4\{\nu_7\circ\Sigma^6 p\}\oplus\mathbb{Z}_3\{\alpha_1(7)\circ\Sigma^6 p\}$.
(5) $G_7(\mathbb{C}P^2, S^8) = 0$.
(6) $G_8(\mathbb{C}P^2, S^9) = \mathbb{Z}_2\{2\nu_9\circ\Sigma^8 p\}\oplus\mathbb{Z}_3\{\alpha_1(9)\circ\Sigma^8 p\}$.

Theorem 5. [16, Corollary 2.4] *For each $n\geq 1$. we have an isomorphism of abelian groups*

$$G_n(map(X,Y;0))\cong G_n(Y)\oplus G(\Sigma^n X, Y).$$

By [16, Corollary 2.4], for each $n, m\geq 1$, we have an isomorphism of abelian groups

$$G_n(map(\Sigma^m\mathbb{C}P^2, S^{m+1};0))\cong G_n(S^{m+1})\oplus G_m(\mathbb{C}P^2, S^{m+1}).$$

Mukai-Golasinski [4] determined Gottelieb groups of sphere $G_{n+k}(S^n)$ for $1\leq k\leq 13$, $2\leq n\leq 26$. Therefore we have group structures of

$$G_n(map(\Sigma^m\mathbb{C}P^2, S^{m+1}:0))\cong G_n(S^{m+1})\oplus G_m(\mathbb{C}P^2, S^{m+1})$$

for $1\leq m\leq 8$ and $1\leq m-n+1\leq 13$.

We recall a long exact sequence [15].

Theorem 6. *There is a long exact sequence*

$$\cdots\to[\Sigma^r A, map(\Sigma B, X; f)]\xrightarrow{w_*}[\Sigma^r A, X]\xrightarrow{P_\alpha}$$

$$\xrightarrow{P_\alpha}[\Sigma(\Sigma^{r-1}A\wedge B), X]\xrightarrow{i'_*}[\Sigma^{r-1}A, map(\Sigma B, X; f)]\to\cdots$$

where $\alpha=[f]\in[\Sigma B, X]$, $P_\alpha(\beta)=[\beta,\alpha]$ is the generalized Whitehead product and $w: map(\Sigma B, X; f)\to B$ is the evaluation map.

From Theorem 4 and 6, we have the following.

Theorem 7. *(1)* $\pi_1(map(\Sigma\mathbb{C}P^2, S^2; f)) \cong \mathbb{Z}_2\{\eta_2 \circ \nu' \circ \Sigma^2 p\} \oplus \mathbb{Z}_3\{[\iota_2, \iota_2] \circ \alpha_1(3)\}$ *for all* $f : \Sigma\mathbb{C}P^2 \to S^2$.

(2) $\pi_2(map(\Sigma^2\mathbb{C}P^2, S^3; f)) \cong \mathbb{Z}_2\{\nu' \circ \overline{2\iota_6}\} \oplus \mathbb{Z}_3\{\overline{\alpha_1(3)}\}$ *for all* $f : \Sigma^2\mathbb{C}P^2 \to S^3$.

(3) (a) For $[f] \in \langle 2\nu_4 \circ \Sigma^3 p, \Sigma\nu' \circ \Sigma^3 p\rangle$, $\pi_3(map(\Sigma^3\mathbb{C}P^2, S^4; f)) \cong \mathbb{Z}_4\{\nu_4^2 \circ \Sigma^6 p\} \oplus \mathbb{Z}_3^2\{\alpha_1(4) \circ \alpha_1(7) \circ \Sigma^6 p, [\iota_4, \iota_4] \circ \alpha_1(7) \circ \Sigma^6 p\}$.

(b) For $[f] \in \{(2n+1)\nu_4 \circ \Sigma^3 p \,|\, n \in \mathbb{Z}\}$ *and* $[g] \in \langle 2\nu_4 \circ \Sigma^3 p, \Sigma\nu' \circ \Sigma^3 p\rangle$, $\pi_3(map(\Sigma^3\mathbb{C}P^2, S^4; f+g)) \cong \mathbb{Z}_2\{\nu_4^2 \circ \Sigma^6 p\} \oplus \mathbb{Z}_3^2\{\alpha_1(4) \circ \alpha_1(7) \circ \Sigma^6 p, [\iota_4, \iota_4] \circ \alpha_1(7) \circ \Sigma^6 p\}$.

(c) For $[f] \in \{\alpha_1(4) \circ \Sigma^3 p, 3\alpha_1(4) \circ \Sigma^3 p\}$ *and* $[g] \in \langle 2\nu_4 \circ \Sigma^3 p, \Sigma\nu' \circ \Sigma^3 p\rangle$, $\pi_3(map(\Sigma^3\mathbb{C}P^2, S^4; f+g)) \cong \mathbb{Z}_4\{\nu_4^2 \circ \Sigma^6 p\} \oplus \mathbb{Z}_3\{\alpha_1(4) \circ \alpha_1(7) \circ \Sigma^6 p\}$.

(4) $\pi_4(map(\Sigma^4\mathbb{C}P^2, S^5; f)) \cong \mathbb{Z}_4\{\overline{\nu_5 \circ \eta_8^2}\} \oplus \mathbb{Z}_3\{\alpha_2(5) \circ \Sigma^8 p\} \oplus \mathbb{Z}_5\{\alpha_{1,5}(5) \circ \Sigma^8 p\}$ *for all* $f : \Sigma^4\mathbb{C}P^2 \to S^5$.

(5) (a) For $[f] \in \langle 2\nu_6 \circ \Sigma^5 p\rangle$, $\pi_5(map(\Sigma^5\mathbb{C}P^2, S^6; f)) \cong \mathbb{Z}_4^2\{\nu_6 \circ g_9, \overline{\nu}_6 \circ \Sigma^{10} p\} \oplus \mathbb{Z}_3\{[\iota_6, \iota_6] \circ \alpha_1(11) \circ \Sigma^{10} p\}$.

(b) For $[f] \in \{\nu_6 \circ \Sigma^5 p, 3\nu_6 \circ \Sigma^5 p\}$ *and* $[g] \in \langle 2\nu_6 \circ \Sigma^5 p\rangle$, $\pi_5(map(\Sigma^5\mathbb{C}P^2, S^6; f)) \cong \mathbb{Z}_4\{\nu_6 \circ g_9\} \oplus \mathbb{Z}_2\{\overline{\nu}_6 \circ \Sigma^{10} p\} \oplus \mathbb{Z}_3\{[\iota_6, \iota_6] \circ \alpha_1(11) \circ \Sigma^{10} p\}$.

(c) For $[f] \in \{\alpha_1(6) \circ \Sigma^5 p, 2\alpha_1(6) \circ \Sigma^5 p\}$ *and* $[g] \in \langle 2\nu_6 \circ \Sigma^5 p\rangle$, $\pi_5(map(\Sigma^5\mathbb{C}P^2, S^6; f)) \cong \mathbb{Z}_4^2\{\nu_6 \circ g_9, \overline{\nu}_6 \circ \Sigma^{10} p\}$.

(6) $\pi_6(map(\Sigma^6\mathbb{C}P^2, S^7; f)) \cong \mathbb{Z}_8\{\sigma' \circ \overline{2\iota_{14}}\} \oplus \mathbb{Z}_3\{\overline{\alpha_2(7)}\} \oplus \mathbb{Z}_5\{\overline{\alpha_{1,5}(7)}\}$ *for all* $f : \Sigma^6 p \to S^7$.

(7) (a) $\pi_7(map(\Sigma^7\mathbb{C}P^2, S^8; *)) \cong \mathbb{Z}_4^2\{\sigma_8 \circ \nu_{15} \circ \Sigma^{14} p, \nu_8 \circ \sigma_{11} \circ \Sigma^{14} p\} \oplus \mathbb{Z}_3^2\{\beta_1(8) \circ \Sigma^{14} p, [\iota_8, \iota_8] \circ \alpha_1(15) \circ \Sigma^{14} p\}$.

(b) For $[f] \in \{\nu_8 \circ \Sigma^7 p, 3\nu_8 \circ \Sigma^7 p\}$, $\pi_7(map(\Sigma^7\mathbb{C}P^2, S^8; f)) \cong \mathbb{Z}_4\{\sigma_8 \circ \nu_{15} \circ \Sigma^{14} p + \nu_8 \circ \sigma_{11} \circ \Sigma^{14} p\} \oplus \mathbb{Z}_3^2\{[\iota_8, \iota_8] \circ \alpha_1(5) \circ \Sigma^{14} p, \beta_1(8) \circ \Sigma^{14} p\}$.

(c) For $[f] = 2\nu_8 \circ \Sigma^7 p$, $\pi_7(map(\Sigma^7\mathbb{C}P^2, S^8; f)) \cong \mathbb{Z}_4\{\sigma_8 \circ \nu_{15} \circ \Sigma^{14} p\} \oplus \mathbb{Z}_2\{\nu_8 \circ \sigma_{11} \circ \Sigma^{14} p\} \oplus \mathbb{Z}_3^2\{\beta_1(8) \circ \Sigma^{14} p, [\iota_8, \iota_8] \circ \alpha_1(15) \circ \Sigma^{14} p\}$.

(d) $[f] \in \{\alpha_1(9)\circ\Sigma^7 p, 2\alpha_1(9)\circ\Sigma^7 p\}$, $\pi_7(map(\Sigma^7\mathbb{C}P^2, S^8; f)) \cong \mathbb{Z}_4^2\{\sigma_8\circ\nu_{15}\circ\Sigma^{14}p, \nu_8\circ\sigma_{11}\circ\Sigma^{14}p\} \oplus \mathbb{Z}_3\{\beta_1(8)\circ\Sigma^{14}p\}$.

(8) (a) For $[f] \in \langle\alpha_1(9)\circ\Sigma^8 p, 2\nu_9\circ\Sigma^8 p\rangle$, $\pi_8(map(\Sigma^8\mathbb{C}P^2, S^9; f)) \cong \mathbb{Z}_8\{\nu_9^2\circ g_{15}\} \oplus \mathbb{Z}_2^2\{\overline{\nu}_9\circ\nu_{17}\circ\Sigma^{16}p, \overline{\eta_9\circ\varepsilon_{10}}\} \oplus \mathbb{Z}_7\{\alpha_{1,7}(9)\circ\Sigma^{14}p\} \oplus \mathbb{Z}_9\{\alpha_3'(9)\circ\Sigma^{14}p\}$.

(b) For $[f] \in \{\nu_9\circ\Sigma^8 p, 3\nu_9\circ\Sigma^8 p\}$ *and* $[g] \in \langle\alpha_1(9)\circ\Sigma^8 p, 2\nu_9\circ\Sigma^8 p\rangle$, $\pi_8(map(\Sigma^8\mathbb{C}P^2, S^9; f+g)) \cong \mathbb{Z}_2\{\nu_9^2\circ g_{15}\} \oplus \mathbb{Z}_2\{\overline{\eta_9\circ\varepsilon_{10}}\} \oplus \mathbb{Z}_7\{\alpha_{1,7}(9)\circ\Sigma^{14}p\} \oplus \mathbb{Z}_9\{\alpha_3'(9)\circ\Sigma^{14}p\}$.

We denote the path-components of mapping space $map(X, Y)$ containing f by $map(f)$.

Finally, we classify homotopy types of path-components of the following mapping spaces.

Theorem 8. *(1)* $map(\Sigma^n\mathbb{C}P^2, S^{n+1}; f)$ *consists of one path-component for* $n = 1, 2, 4, 6$.

(2) $map(\Sigma^3\mathbb{C}P^2, S^4; f)$ *consists of four path-components* $map(f)$, $map(f+g)$, $map(f+h)$ *and* $map(f+g+h)$ *where* $[f] \in \langle 2\nu_4\circ\Sigma^3 p, \Sigma\nu'\circ\Sigma^3 p\rangle$, $[g] \in \{(2n+1)\nu_4\circ\Sigma^3 p \mid n \in \mathbb{Z}\}$ *and* $[h] \in \{\alpha_1(4)\circ\Sigma^3 p, 2\alpha_1(4)\circ\Sigma^3 p\}$.

(3) $map(\Sigma^5\mathbb{C}P^2, S^6; f)$ *consists of four path-components* $map(f)$, $map(f+g)$, $map(f+h)$ *and* $map(f+g+h)$ *where* $[f] \in \langle 2\nu_6\circ\Sigma^5 p\rangle$, $[g] \in \{\nu_6\circ\Sigma^5 p, 3\nu_6\circ\Sigma^5 p\}$ *and* $[h] \in \{\alpha_1(6)\circ\Sigma^5 p, 2\alpha_1(6)\circ\Sigma^5 p\}$.

(4) $map(\Sigma^7\mathbb{C}P^2, S^8; f)$ *consists of six path-components* $map(*)$, $map(f)$, $map(2f)$, $map(g)$, $map(f+g)$ *and* $map(2f+g)$ *where* $[f] = \nu_8\circ\Sigma^7 p$ *and* $[g] = \alpha_1(8)\circ\Sigma^7 p$.

(5) $map(\Sigma^8\mathbb{C}P^2, S^9; f)$ *consists of two path-components* $map(f)$ *and* $map(f+g)$ *where* $[f] = \nu_8\circ\Sigma^7 p$ *and* $[g] = \alpha_1(8)\circ\Sigma^7 p$.

References

[1] M. Arkowitz, The generalized Whitehead product, Pacific J. Math., **12** , (1962), 7–23. MR 27 ♯5262 Zbl 0118.18404

[2] M. G. Barratt, P. J. Hilton, On join operations in homotopy groups, Proc. London Math. Soc. (3), **3**, (1953), 430–445. MR 15,643a Zbl 0053.43403

[3] J. B. Gatsinzi, Rational Gottlieb group of function spaces of maps into an even sphere, Int. J. Algebra, **6**, (2012), no. 9–12, 427–432. MR 2942988 Zbl 1252.55007

[4] M. Golasiński, J. Mukai, Gottlieb groups of spheres, Topology, **47**, (2008), no. 6, 399–430. MR 2009j:55017 Zbl 1172.55004

[5] M. Golasiński, J. Mukai, Gottlieb and Whitehead center groups of spheres, projective and Moore spaces, MR 3243535 Zbl 1312.55001

[6] D. H. Gottlieb, Evaluation subgroups of homotopy groups, Amer. J. Math., **91**, (1969), 729–756. MR 43 ♯1181 Zbl 0185.27102

[7] B. Gray, Homotopy theory. An introduction to algebraic topology, Pure and Applied Math., Vol. **64**, Academic Press, New York-London, (1975). MR 53 ♯6528 Zbl 0322.55001

[8] V. L. Hansen, Equivalence of evaluation fibrations, Invent. Math., **23**, (1974), 163–171. MR 51 ♯4242 Zbl 0281.55002

[9] D. Harris, Every space is a path component space, Pacific J. Math., **91**, (1980), no. 1, 95–104. MR 82d:54006 Zbl 0468.54005

[10] P. J. Hilton, On the homotopy groups of the union of spheres, J. Lond. Math. Soc, **30**, (1955), 154–172. MR 16,847d Zbl 0064.17301

[11] H. Kachi, J. Mukai, T. Nozaki, Y. Sumita, D. Tamaki, Some cohomotopy groups of suspended projective planes, Math. J. Okayama Univ., **43**, (2001), 105–121. MR 2003e:55016 Zbl 1039.55010

[12] P. J. Kahn, Some function spaces of CW type, Proc. Amer. Math. Soc., **90**, (1984), no. 4, 599–607. MR 85j:55027 Zbl 0511.57013

[13] L. Kristensen, I. Madsen, Note on Whitehead products in spheres, Math. Scand., **21**, (1967), 301–314, (1969). MR 39 ♯6316 Zbl 0176.21002

[14] S. Kikkawa, J. Mukai, D.Takaba, Cohomotopy sets of projective planes, J. Fac. Sci. Shinshu Univ., **33**, (1998), no. 1, 1–7. MR 99j:55010 Zbl 0922.55009

[15] G. E. Lang Jr, The evaluation map and EHP sequences, Pacific J. Math., **44**, (1973), 201–210. MR 49 ♯6235 Zbl 0248.55016

[16] G. Lupton, S. B. Smith, Gottlieb Groups of Function Spaces, arXiv:1302.1826v1.

[17] G. Lupton, S. B. Smith, Criteria for components of a function space to be homotopy equivalent, Math. Proc. Cambridge Philos. Soc., **145**, (2008), no. 1, 95–106. MR 2009e:55009 Zbl 1151.55006

[18] K-I. Maruyama, H. Oshima, Homotopy groups of the spaces of self-maps of Lie groups, J. Math. Soc. Japan, **60**, (2008), no. 3, 767–792. MR 2009h:55014 Zbl 1148.55002

[19] J. P. May, A concise course in algebraic topology, Chicago Lectures in Math., University of Chicago Press, (1999). MR 2000h:55002 Zbl 0923.55001

[20] C. A. McGibbon, Self-maps of projective spaces, Trans. Amer. Math. Soc., **271**, (1982), no. 1, 325–346. MR 83h:55007 Zbl 0491.55014

[21] J. Milnor, On spaces having the homotopy type of a CW-complex, Trans. Amer. Math. Soc., **90**, (1959), 272–280. MR 20 ♯6700 Zbl 0084.39002

[22] M. Mimura, G. Nishida, H. Toda, Localization of CW-complexes and its applications, J. Math. Soc. Japan, Vol. **23**, (1971), No. 4, 593–624. MR 45 ♯4413 Zbl 0217.48801

[23] J. Mukai, On stable homotopy of the complex projective space, Japan J. of Math., **19**, (1993), 191–216. MR 95c:55014 Zbl 0810.55006

[24] J. Mukai, Note on existence of the unstable Adams map, Kyushu J. Math., **49**, (1995), no. 2, 271—279. MR 98e:55015 Zbl 0855.55012

[25] K. Oguchi, Generators of 2-primary components of homotopy groups of spheres, unitary groups and symplectic groups, J. Fac. Sci. Univ. Tokyo Sect. I, **11**, (1964), 65–111, (1964). MR 30 ♯4262 Zbl 0129.38902

[26] S. B. Smith, Rational homotopy of the space of self-maps of complexes with finitely many homotopy groups, Trans. Amer. Math. Soc., **342**, (1994), no. 2, 895–915. MR 94j:55012 Zbl 0802.55009

[27] N. E. Steenrod, A convenient category of topological spaces, Michigan Math. J., **14**, (1967), 133–152. MR 35 ♯970 Zbl 0145.43002

[28] H. Toda, Composition methods in homotopy groups of spheres, Ann. of Math. Studies, **49**, Princeton, (1962). MR 26 ♯777 Zbl 0101.40703

[29] K. Varadarajan, Generalised Gottlieb groups, J. Indian Math. Soc., (N.S.), **33**, (1969), 141–164. MR 43 ♯6926 Zbl 0199.26501

[30] G. W. Whitehead, On products in homotopy groups, Ann. of Math. (2), **47**, (1946), 460–475. MR 8,50b Zbl 0060.41106

Jin Ho Lee
Department of Mathematics
Korea University
Seoul 136-701, Korea
E-mail address: `sabforev@korea.ac.kr`